

AGDOXYTYL - water soluble antibiotic powder



Composition:

| |
|----------------------------|
| Each gram powder contains: |
| Tylosine tartrate. 20 gm |
| Doxycycline Hcl. 15 gm |

Indications:

This combination of two antibiotics is active against many Gram-positive and Gram-negative bacteria, as well as against mycoplasmas.

To be used for the treatment of infections caused by Gram-positive and Gram-negative bacteria, susceptible to tylosin and/or doxycycline in poultry, calves and pigs, especially infections of the respiratory tract and gastro-intestinal tract.

Contra-indications: Hypersensitivity to tetracyclines, serious renal- and/or liver dysfunctions. Do not administer simultaneously with penicillins and/or cephalosporines.

Dosage and Administration: Administer orally via drinking water or calf-milk.
Poultry: Preventive dose: 100 g per 300 litres of drinking water, during 3-5 days.
Therapeutic dose : 100 - 200 g per 150 litres of drinking water, during 3-5 days.
Calves: 100 g per 80 calves of 50 kg bodyweight, during 3-5 days.
Pigs: 100 g per 200 litres of drinking water, during 3-5 days.

Recommendations and warnings: Administer during treatment only medicated water. Medicated drinking water should be used within 24 hours after preparation.

Withdrawal time: For meat: 15 days before slaughter. For eggs: 10 days.

Storage: Keep closed in a cool and dry place, away from sunlight and safe from children.

Packing:

12 x 10 x 100 gm sachet in 1 carton.
12 x 1 kg jar in 1 carton.

FOR VETERINARY USE ONLY